

Pacific University

CommonKnowledge

College of Optometry

Theses, Dissertations and Capstone Projects

1979

The estimation of corneal rigidity by means of comparative tonometry

Timothy G. Moore
Pacific University

Barry A. Eichenbaum
Pacific University

Recommended Citation

Moore, Timothy G. and Eichenbaum, Barry A., "The estimation of corneal rigidity by means of comparative tonometry" (1979). *College of Optometry*. 536.
<https://commons.pacificu.edu/opt/536>

This Thesis is brought to you for free and open access by the Theses, Dissertations and Capstone Projects at CommonKnowledge. It has been accepted for inclusion in College of Optometry by an authorized administrator of CommonKnowledge. For more information, please contact CommonKnowledge@pacificu.edu.

The estimation of corneal rigidity by means of comparative tonometry

Abstract

The estimation of corneal rigidity by means of comparative tonometry

Degree Type

Thesis

Degree Name

Master of Science in Vision Science

Committee Chair

Lynn J. Coon

Subject Categories

Optometry

Copyright and terms of use

If you have downloaded this document directly from the web or from CommonKnowledge, see the "Rights" section on the previous page for the terms of use.

If you have received this document through an interlibrary loan/document delivery service, the following terms of use apply:

Copyright in this work is held by the author(s). You may download or print any portion of this document for personal use only, or for any use that is allowed by fair use (Title 17, §107 U.S.C.). Except for personal or fair use, you or your borrowing library may not reproduce, remix, republish, post, transmit, or distribute this document, or any portion thereof, without the permission of the copyright owner. [Note: If this document is licensed under a Creative Commons license (see "Rights" on the previous page) which allows broader usage rights, your use is governed by the terms of that license.]

Inquiries regarding further use of these materials should be addressed to: CommonKnowledge Rights, Pacific University Library, 2043 College Way, Forest Grove, OR 97116, (503) 352-7209. Email inquiries may be directed to: copyright@pacificu.edu

The Estimation of Corneal Rigidity
by Means of
Comparative Tenometry

Investigators: Timothy G. Moore and Barry A. Eichenbaum


Faculty Advisor: Dr. Lynn J. Coon, O.D.

Submitted in partial fulfillment of the requirements for the
degree of Doctor of Optometry at Pacific University College
of Optometry.

TABLE OF CONTENTS


Thesis Acceptance.	I
Acknowledgments.	II
Sample Subject Release Form.	III
Review of the Literature.	1-6
Methodology.	7-8
Results and Discussion.	9-11
Summary.	11-13
Table I.	14
Figures I-IX.	15-23
Appendix.	24-26
Bibliography.	27-33


Thesis submitted by:


Timothy G. Moore


Barry A. Eichenbaum

Accepted by:


Lynn J. Coen, O.D.


Grade

This study was funded by a grant from Beta Sigma Kappa Optometric Fraternity. The authors also wish to thank Lynn J. Coon, O.D. for his guidance and assistance in this undertaking.

III

SUBJECT RELEASE FORM

I. Basic Information

Project Title: The Estimation of Corneal Rigidity by
Means of Comparative Tonometry
Investigators: Timothy G. Moore and Barry A. Eichenbaum
Advisor: Lynn J. Coon, O.D.
Location: Pacific University College of Optometry,
Forest Grove, Oregon and Portland Optometric Clinic,
Portland, Oregon.
Date: Fall 1978

II. Description of Project

The intent of this project is to attempt to estimate corneal rigidity using a number of subjects. This will be done by comparing the readings obtained by two forms of tonometry (a reading commonly taken to aid in determining the presence of glaucoma). Measurements will be taken by two standard forms of tonometry, the Schiottz Method and the Non-contact or air-puff Method. The Schiottz Method does involve actually touching the cornea and a local anesthetic in the form of eye drops will be administered to avoid any subject discomfort. The cornea will be examined before and after the tonometry.

III. Description of Risks

Risks are very minimal. Some subjects do report some irritation following tonometry. Some slight corneal abrasion may occur but is not likely. Subjects will be thoroughly checked before they are dismissed.

IV. Description of Benefits

Subjects will have a more precise knowledge of their intraocular pressure and be informed as to the status of their cornea's health. Corneal rigidity may be important in contact lens fitting for the purpose of myopia (near-sightedness) reduction.

V. Offer to Answer Any Inquiries

The experimenters will be more than happy to answer any questions you may have during the course of the project.

VI. Time Required

The subject can plan on donating no more than thirty minutes of their time to this project. This will be done in one visit.

VII. Freedom to Withdraw

You are free to withdraw your consent and discontinue your participation in this project at any time without prejudice to you.

I have read and understand the above.
I am 18 years of age or older.

Signed _____ date _____

Review of the Literature

Tonometry is defined as "the measurement of ocular tension with a tonometer" (Schapero et.al. 1968), which still leaves one in doubt as to just what tonometry is, unless of course we define ocular tension. Ocular tension is in turn defined as "the resistance of the tunics of the eye to indentation, which depends upon the intraocular pressure, the thickness and rigidity of the tunics, the surface area, etc.,....." (Schapero et.al. 1968). Ideally one would wish to be able to directly measure the pressure within the eye by a means which would eliminate intervening factors such as rigidity of the tunics. Cannulation of the eye so that the internal pressure within the eye is directly measured by a mercury manometer (Kaufman 1972) would avoid the problems of mechanical indirect measurements which are influenced by technical difficulties as well as resistance to mechanical deformation or indentation by the elasticity of the tunics of the eye. Although it is perhaps the best theoretical means of determining intraocular pressure, cannulation presents many problems of its own. Anesthetics necessary for cannulation would most likely be a general anesthetic, usually barbiturates. Barbiturates have been found to lower the intraocular pressure (Moses 1975). Leakage of the aqueous and trauma to the eye are also factors which rule out cannulation as a practical means of determining ocular tension. Alternatively we are left with several methods of indirectly determining intraocular pressure by mechanical devices.

Two major approaches for the determination of ocular tension are currently in use; indentation tonometry and applanation tonometry, each with their inherent drawbacks and advantages. Indentation or Schiottz tonometry was for many years considered to be the standard method of determining intraocular pressure. More recently with the development of the Goldmann applanation tonometer, which is generally considered more accurate and more reliable, (Bengtsson 1972B, Moses 1958, Smith 1964, Kaufman 1972) has replaced the Schiottz indentation method as the standard.

The Schiottz tonometer is subject to a great degree of variability even when used by skilled persons (Glaster 1965). Applanation tonometry has been found to be subject to less variability than Schiottz tonometry (Bengtsson 1972B). Scott(1971) using several tonometers (Goldmann, Mackay-Marg, Maklakov and Schiottz) found the highest correlation between Schiottz and Goldmann tonometric findings. Population means for Goldmann applanation tonometry and Schiottz indentation tonometry have been found to have a high correlation although individual findings do not have such a high correlation (Armaly 1960).

Disparity between applanation and indentation tonometry has been to a large extent attributed to ocular rigidity (Buxton 1971, Moses 1958). Much of this disparity is attributed to the errors inherent in indentation tonometry. Factors which have been found to influence ocular rigidity are as follows: (from Buxton 1971)

A. Decreased ocular rigidity

1. Water provocative test
2. Endocrine exophthalmos
3. Myopia
4. Strong miotics in glaucoma therapy
5. Buphthalmic eyes
6. Megalocornea
7. Intraocular surgery
8. Hydration of the cornea

B. Increased ocular rigidity

1. Small cornea
2. Increased corneal thickness
3. Contraction of extraocular muscles

C. Questionable effects on ocular rigidity

1. Increased age
2. Escape of intraocular fluid by instrumentation

Factors which affect intraocular pressure more so than ocular rigidity would tend to have the same effect on indentation and applanation tonometry so that they would have minimal effects in the determination of ocular rigidity. These factors include blood pressure, sex, season of the year and time of day, all of which were considered to have independent effects on intraocular pressure (Bengtsson 1972A).

Applanation tonometry, more specifically the Goldmann method, is considered to be devoid of the influences of ocular rigidity and is thereby considered to be a more direct measure of actual intraocular pressure (Borish 1975).

When a surface of three to four millimeters is applanated the surface tension due to tears and acting in a direction opposite to the force of resistance due to ocular rigidity equals and cancels the force of ocular rigidity (Moses 1975).

The following formula is used as a basis for applanation tonometry; the Imbert-Fick Law:

$$P=F/A,$$

where P=pressure, F force to applanate an area, A. A modification of this formula,

$$P=\frac{F+M-N}{A}$$

includes correction for force due to tear surface tension, M, and the force of resistance to corneal deformation, N (Moses 1975).

Formulations involving Schiotz tonometry are based primarily on the experimental data and work of Friedenwald (1937, 1954, 1957). From Borish (1975) Friedenwald's Formula for Schiotz tonometry is as follows:

$$a + bS = \frac{W \text{ (weight of the total plunger - indenting force)}}{P_t \text{ (the IOP with the tonometer on the eye)}}$$

where a and b are experimentally determined constants and S is the Schiotz scale reading. Clinically such a formulation is impractical and a nomogram devised by Friedenwald on a semilog plot is used for determination of intraocular pressure as well as an ocular rigidity coefficient (Borish 1975).

The difference between tonometric findings using the Schiotz tonometer and the Goldmann tonometer have generally been attributed to ocular rigidity as stated previously, since ocular rigidity is considered to have no effect on applanation tonometry. Regression analysis of data for Schiotz and Goldmann tonometry indicate a deviation from each other above and below the means and have been referred to as a regression effect (Levene 1971). Bengtsson (1973) suggests that

regression curves deviating from Friedenwald's 1955 calibration curve are a consequence of many kinds of disagreement between the two methods of measurement.

Use of a differential in determination of ocular rigidity is not without precedent. Several values taken with the Schiotz tonometer using different calibration weights have been used on Friedenwald's nomogram in determination of a coefficient of ocular rigidity. Syrdalen (1970) used a Goldmann-Schiotz differential in formulating a means for determination of an ocular rigidity coefficient as follows:

$$K = \frac{\log P_t - \log(\text{appl. ton.} + 1)}{V_c - 0.44 \text{ mm}^3},$$

where K = the rigidity coefficient, P_t = intraocular pressure during tonometry (Friedenwald's table 1955), V_c = volume of corneal indentation caused by Schiotz tonometry (Friedenwald 1955), 0.44 mm^3 = corneal indentation caused by applanation tonometry (Goldmann and Schmidt 1957). The addition of 1mm to the applanation tonometry values is to compensate for the increase in intraocular pressure from sitting to reclining as also found by Scott (1971).

The advent of modern technology has not passed up the area of tonometry. Many new and sophisticated electronic devices for measurement of intraocular pressure have appeared on the market in the past few years. Some of these new devices are no more reliable than previous instrumentation, specifically the Goldmann tonometer, which remains the standard by which the others are compared and calibrated. One of the most elaborate of these new devices is the

American Optical Non-Contact Tonometer. The A.O. Non-Contact tonometer is an applanation tonometer which applanates the cornea by means of compressed air. Many factors which may affect the reliability of readings with the Goldmann tonometer are circumvented by use of the A.O. Non-Contact tonometer. Factors introducing errors into Goldmann tonometric measurements are; surface tension of the anesthetic, improperly cleaned prism, improperly adjusted height, prolonged contact with the cornea, corneal edema, improper light placement, width of the fluorescein meniscus (Moses 1958). When properly calibrated and used as prescribed, the A.O. Non-Contact tonometer is not limited by any of these factors. Tests for reliability and effect on corneal integrity (Forbes et. al. 1974) show a correlation coefficient of +0.9 for the A.O. Non-Contact tonometer compared to the Goldmann tonometer. In this same study, repeated readings with the A.O. NCT showed no insult to the corneal epithelium.

The advantages and ease of use over Goldmann applanation tonometry are major factors in selection of the A.O. Non-Contact tonometer for this study. Applanation by means of the A.O. NCT will be compared with differential Schiötz readings in an attempt to ascertain a relationship between the differential of comparative tonometry and an ocular rigidity coefficient.

Methodology

Subjects for this study are to be selected from the general population. The only restriction upon selection of subjects will be that the subject's corneas have never been subject to any trauma which resulted in permanent corneal defects which are visible upon examination with the biomicroscope.

Prior to testing, each subject will be required to sign a release form describing the experiment and possible risks which may be involved.

Subjects will be examined with a biomicroscope before any tonometric measurements are taken to determine the state of corneal integrity. Pre-tonometric examination with the biomicroscope will also help to provide a basis for comparison of the corneal integrity with a second post-tonometric biomicroscopic examination to ascertain the extent of corneal damage due to tonometric procedures. The second biomicroscopic exam will be performed with sodium fluorescein to highlight any epithelial damage.

The actual experimental procedure will involve taking standard tonometric readings with the American Optical Non-Contact applanation tonometer and readings with the Schiottz indentation tonometer. Measurements of intraocular pressure will be made first with the Non-Contact tonometer to avoid any possible "massaging" effects due to volumetric displacement which occurs with Schiottz tonometry. A minimum of three

"valid" measurements of intraocular pressure will be made of each eye with the non-contact tonometer with an average of the three taken as the intraocular pressure.

Tonometric measurements will be taken with the Schiotz tonometer after administration of the appropriate local corneal anesthetic. Measurements will be taken in the standard fashion with the subject reclining. Three measurements will be made upon each eye. The three measurements will be made with three different calibration weights so that a differential comparison may be made as in determination of the Friedenwald coefficient of ocular rigidity. Comparisons will then be made between the differential Schiotz measurements and those obtained with the American Optical Non-Contact tonometer. Appropriate methods of statistical analysis will be employed.

Results and Discussion

The premise of the method of comparative tonometry is based on the idea that tonometric measurements of a subject taken by two different methods should differ from the mean of a population by the same amount if the two methods of measurement were linearly related. If this were the case, one would expect that a ratio of the difference from the mean divided by the mean would be equal in the two methods, i.e.,

$$\frac{\bar{X}_{\text{NCT}} - \text{NCT Reading}_n}{\bar{X}_{\text{NCT}}} = \frac{\bar{X}_{\text{Schietz}} - \text{Schietz Reading}_n}{\bar{X}_{\text{Schietz}}}$$

The ratios were assigned a negative value if the subject's reading was below the population mean and a positive value if the reading was above population mean. Differences between the ratios obtained from the NCT and those obtained from the Schietz were taken as a measure of non-linearity between the two methods of tonometry.

It was our contention that these differences in values were related to corneal rigidity. The term, "corneal rigidity", is used somewhat loosely since comparisons were made using Friedenwald's Coefficient of Ocular Rigidity. To our knowledge, no comparisons have been made between readings taken at points on non-corneal surfaces and corneal surfaces. Technically, in this study we can not distinguish between ocular and corneal rigidity although the primary mechanical resistance to deformation is at the cornea when Schietz

tonometry is used. Hydrestatic resistance is presumed here to remain constant. The mechanical resistance of the cornea to greater or lesser degrees of deformation, dependent upon the Schiøtz tonometer plunger load, is taken to be due to corneal rigidity.

A coefficient of ocular rigidity was determined for each subject by graphical projection on Friedenwald's nemi-gram (taken from Adler, 1975). The coefficient was obtained by using an average of three Non-contact tonometer readings and three Schiøtz readings, each using a different plunger load. These values may be found in Table I.

In determining a measure of rigidity from comparative tonometry, the ratios obtained as previously described, may be both positive, both negative or one positive and the other negative. What was of primary concern was the difference between the NCT ratio and the Schiøtz ratio,

$$\frac{\bar{X}_{NCT} - NCT_n}{\bar{X}_{NCT}} - \frac{\bar{X}_{Schiøtz} - Schiøtz_n}{\bar{X}_{Schiøtz}}$$

The difference between the two ratios was taken as a measure of corneal rigidity. This may be better visualized in diagrams I, II, and III. The higher the rigidity the higher the positive value. Values giving no difference can not be construed to have no rigidity but are points where there is a linear relationship between NCT and Schiøtz tonometry. Values above this 1:1 relationship have a higher rigidity

and those below it are of a lower rigidity. It is this non-linearity between the two different methods of tonometry that forms the basis for the quantification of corneal rigidity in this study.

The determination of corneal rigidity is based on population means for Non-contact and Schiøtz tonometry done in this study. (See figures IV, V, VI, VII, and VIII.) The sample size of this study was somewhat small, but was adequate for statistical analysis. The Pearson product-moment correlation coefficient was used to compare figures obtained with comparative tonometry with those obtained from Friedenwald's nomogram, which is shown as figure IX. Correlation between the Ocular Rigidity Coefficient obtained from the Friedenwald nomogram and the NCT - 5.5gr. Schiøtz values was only .42. Correlations with the NCT - 7.5gr. and NCT - 10gr. pairs to the Friedenwald coefficients were somewhat higher; .52 and .69 respectively. These last two comparisons show a fairly high positive correlation with the values of the Friedenwald nomogram.

Summary

It would be injudicious at this time to state that the method of assessing corneal rigidity as described in this study is valid beyond any doubt. Several aspects of the study limit the reliability of this method as an indicator

of corneal rigidity. First of all, it is dependent upon population means, which in this case are based on a rather small population. Frequency distributions are illustrated superimposed on theoretical normal distributions in figures IV, V, VI, and VII. A comparison of theoretical normal distributions for the NCT, 5.5gr., 7.5gr. and 10.0gr. Schiotz findings of the population used in this study are illustrated in figure VIII. Here, it may be seen that there is a considerable difference in the distribution of the 5.5gr. Schiotz readings. This may or may not have had an effect on the correlation coefficient between the NCT - 5.5gr. Schiotz comparative rigidity measure and the Friedenwald Ocular Rigidity Coefficient. A study in which a larger population is used will probably provide more conclusive information.

The usefulness of an ocular rigidity coefficient has been considered to be of dubious value in clinical practice. (Adler, 1975). An ancillary aspect of this study may add credence to that statement. In an attempt to relate corneal (ocular) rigidity and Orthokeratology, a comparison between rigidity and acuity improvement was made with several patients involved in the Orthokeratology study being conducted at Pacific University College of Optometry. Correlation between the increase in decimal acuity and Friedenwald's coefficient of ocular rigidity was very low, .08 to be exact.


This study could provide the basis for a more comprehensive study in which the reliability of the comparative method may be more conclusively determined.

TABLE I

Friedenwald's Ocular Rigidity Coefficient
as Determined by
Graphical Plot of Schiøtz and NCT Readings


<u>Subject No.</u>	<u>Friedenwald's Ocular Rigidity Coefficient</u>
1	.016
2	.019
3	.021
4	.017
5	.019
6	.024
7	.019
8	.018
9	.020
10	.022
11	.026
12	.020
13	.0225
14	.022
15	.0175
16	.018
17	.021
18	.0175
19	.0195
20	.021
21	.020
22	.023
23	.0195
24	.0205
25	.0305
26	.0175
27	.0205
28	.022

FIGURE I




Rigidity values obtained by comparative method, using the Non-contact and 5.5gr. Schiøtz tonometers.

FIGURE II





Rigidity values obtained by comparative method, using the Non-contact and 7.5gr. Schiötz tonometers.

FIGURE III





Rigidity values obtained by comparative method, using the Non-contact and 10.0gr. Schiotz tonometers.

FIGURE IV




Frequency distribution of readings with the Non-contact
tenometer superimposed on the theoretical normal distribution
for the population.

FIGURE V





Frequency distribution of readings with the 5.5gr. Schietz tonometer superimposed on the theoretical normal distribution for the population.

FIGURE VI




Frequency distribution of readings with the 7.5gr. Schiøtz tonometer superimposed on the theoretical normal distribution for the population.

FIGURE VII



Frequency distribution of readings with the 10.0gr. Schiøtz tonometer superimposed on the theoretical normal distribution for the population.


FIGURE VIII



Open circle = Non-contact tonometer distribution
Closed circle = 5.5gr. Schiøtz tonometer distribution
Square = 7.5gr. Schiøtz tonometer distribution
Triangle = 10.0gr. Schiøtz tonometer distribution

Theoretical normal distributions of Non-contact, 5.5gr.,
7.5gr., and 10.0gr. Schiøtz tonometry values for the
experimental population.

FIGURE IX



Friedenwald nomogram taken from Adler's Physiology
of the Eye, 1975.

APPENDIX I

Mean Values and Deviation from the Mean

	<u>N.C.T.</u>	<u>5.5gr.Schietz</u>	<u>7.5gr.Schietz</u>	<u>10.0gr.Schietz</u>
Mean Value(mm Hg)	13.57	14.96	12.61	11.79

	<u>Deviation from the Mean (mm Hg)</u>			
<u>Subject No.</u>	<u>N.C.T.</u>	<u>5.5gr.Schietz</u>	<u>7.5gr.Schietz</u>	<u>10.0gr.Schietz</u>
1	-3.9	2.04	1.39	4.21
2	-1.57	0.04	-1.61	-0.79
3	1.76	2.04	1.39	1.21
4	2.43	4.04	6.39	4.21
5	-0.9	-4.96	-2.61	-0.79
6	6.1	1.04	1.39	2.21
7	-0.9	-1.96	-0.61	-0.79
8	-3.9	6.04	0.39	-0.79
9	-0.57	-1.96	-1.61	-0.79
10	1.76	0.04	-0.61	-0.79
11	-1.9	-4.96	-3.61	-6.79
12	-0.57	1.04	3.39	-0.79
13	1.76	-2.96	-1.61	-1.79
14	-0.57	-2.96	-1.61	-1.79
15	-2.24	2.04	0.39	2.21
16	-2.57	-2.96	-1.61	-0.79
17	-2.43	2.04	0.39	2.21
18	-2.43	6.04	6.39	4.21
19	-2.57	-4.96	-4.61	-1.79
20	6.1	1.04	3.39	2.21
21	0.43	0.04	0.39	1.21
22	-3.24	4.04	0.39	1.21
23	0.9	-1.96	-1.61	-0.79
24	2.43	1.04	-0.61	1.21
25	-1.24	-2.96	-3.61	-6.79
26	-0.57	6.04	3.39	2.21
27	0.76	-2.96	-0.61	-0.79
28	-0.24	-2.96	-2.61	-1.79

APPENDIX II

Value of the ratio: Deviation from mean(mm Hg.)
Mean(mm Hg.)

<u>Subject No.</u>	<u>N.C.T.</u>	<u>5.5gr.Schietz</u>	<u>7.5gr.Schietz</u>	<u>10.0gr.Schietz</u>
1	-.287	.136	.110	.357
2	-.161	.003	-.128	-.067
3	.130	.136	.110	.103
4	.180	.270	.507	.357
5	-.066	-.332	-.207	-.067
6	.450	.070	.110	.187
7	-.066	-.131	-.048	-.067
8	-.287	.404	.031	-.067
9	-.042	-.131	-.128	-.067
10	.130	.003	-.048	-.067
11	-.140	-.332	-.286	-.576
12	-.042	.070	.269	-.067
13	.130	-.198	-.108	.152
14	-.042	-.198	-.128	.152
15	-.165	.136	.031	.187
16	-.189	-.198	-.128	-.067
17	-.180	.136	.031	.187
18	-.180	.404	.507	.357
19	-.190	-.332	-.366	-.152
20	.450	.070	.269	.187
21	.030	.003	.031	.103
22	-.239	.270	.031	.103
23	-.066	-.131	-.128	-.067
24	.179	.070	-.048	.103
25	-.090	-.198	-.286	-.576
26	-.040	.404	.268	.187
27	.056	-.198	-.048	-.067
28	-.018	-.198	-.207	-.152

APPENDIX III

Value of Rigidity as Determined
by the Equation:

$$\frac{\bar{X}_{NCT} - NCT_n}{\bar{X}_{NCT}} - \frac{\bar{X}_{Schietz} - Schietz_n}{\bar{X}_{Schietz}} = \text{Comparative Rigidity}$$

<u>Subject No.</u>	<u>NCT- 5.5gr.Schietz</u>	<u>NCT-7.5gr.Schietz</u>	<u>NCT-10.0gr.Schietz</u>
1	-.423	-.397	-.644
2	-.119	.012	-.049
3	-.006	.020	.027
4	-.090	-.327	-.177
5	.266	.141	.001
6	.380	.340	.263
7	.065	-.018	.001
8	-.691	-.318	-.220
9	-.089	.086	.025
10	.127	.178	.197
11	.192	.146	.436
12	-.112	-.311	.025
13	.328	.258	-.022
14	.156	.086	-.194
15	-.301	-.196	-.352
16	.009	-.061	-.122
17	-.316	-.211	-.367
18	-.584	-.687	-.537
19	.142	.176	-.038
20	.380	.181	.263
21	.027	-.001	-.073
22	-.509	-.270	-.342
23	.065	.062	.001
24	.109	.227	.076
25	.108	.196	.486
26	-.444	-.308	-.227
27	.254	.104	.123
28	.180	.189	.134

BIBLIOGRAPHY

- Adler, F.H.. Adler's Physiology of the Eye. Sixth Edition.
Chapter 6, "Intraocular Pressure." The C.V. Mosby Co..
St. Louis, Missouri, 1975.
- Andersen, D.R. and W.M. Grant. "Re-evaluation of the Schiøtz
Tonometer Calibration." Investigative Ophthalmology.
Vol. 9: pp. 430-446.
- Aquino, Mario V., Jose T.D. Mayo, Rosie Reyes-Noche and
Arturo Librea. "Ocular Rigidity Among Filipinos."
Philippine Journal of Ophthalmology. Vol. 3(3):
pp. 104-110. July-September, 1971.
- Armaly, M.F.. "Schiøtz Tonometer Calibration and Applanation
Tonometry." Archives of Ophthalmology. Vol. 64: pp. 426-
432. 1960.
- Bengtsson, Bo. "Some Factors Affecting the Distribution of
Intraocular Pressures in a Population." Acta Ophthal-
mologica. (Kbh.) Vol. 50: pp. 33-46. 1972A.
- Bengtsson, Bo. "Comparison of Schiøtz and Goldmann Tonometry
in a Population." Acta Ophthalmologica. Vol. 50(4):
pp. 445-457. 1972B.
- Bengtsson, Bo. "Some Factors Affecting the Relationship
Between Schiøtz and Goldmann Tonometry in a Population."
Acta Ophthalmologica. Vol. 51(6): pp. 798-810. 1973
- Borish, Irving M.. Clinical Refraction. Third Edition.
Chapter 12, "Tonometry." The Professional Press, Inc..
Chicago, 1975.
- Buxton, Jorge N., R.J. Riechers and S.D. Aaron. "Corneal
Grafts and Their Effect Upon the Applanation-Schiøtz
Disparity." Archives of Ophthalmology. Vol. 86(1): pp. 28-
30. July, 1971.

- Dittmar, P., M. Wittenburg, and O. Liegl. "Erfahrungen mit dem Non-Contact Tonometer (Experiences with the Non-Contact Tonometer)." Klinische Monatsblätter für Augenärztliche Fortbildung. Vol. 167(1): pp. 137-141. July, 1975.
- Drance, S. M. "The Coefficient of Scleral Rigidity in Normal and Glaucomatous Eyes." Archives of Ophthalmology Vol. 63: pp. 668-674. 1960.
- Forbes, M., G. Pico and B. Grolman. "A Non-Contact Applanation Tonometer." Archives of Ophthalmology. Vol. 91: pp. 134-140. February, 1974.
- Francois, J., P. Vancea and R Vandekerckhove. "Calibration of Tonometers and True I.O.P." Ophthalmologica (Basel). Vol. 167: pp. 27-48. 1973.
- Friedenwald, J. S.. "Contribution to the Theory and Practice of Tonometry." American Journal of Ophthalmology. Vol. 20: pp. 985-1024. 1937.
- Friedenwald, J. S.. "Calibration of Tonometers." Decennial Report by the Committee on Standardization of Tonometers. American Academy of Ophthalmology. Chapter 7. 1954.
- Friedenwald, J.S.. "Tonometer Calibration: An Attempt to Remove Discrepancies Found in the 1954 Calibration Scale for Schiotz Tonometers." Transcripts of the American Academy of Ophthalmology and Otolaryngology. Vol. 61: p. 108. 1957.
- Gloster, J.. "Tonometry and Tonography." International Ophthalmology Clinic. Vol. 5: pp. 968-1076. 1965.

- Goldmann, H. and Thomas Schmidt. "Über Applanations Tonometrie." *Ophthalmologica*. Vol. 134: pp. 221-242. 1957.
- Hetland-Eriksen, J.. "On Tonometry. 6 Comparative Tonometry with the Goldmann Applanation Tonometer and the Schiotz Tonometer in the Lying Position." *Acta Ophthalmologica*. Vol. 44: p. 522. 1966.
- Kaiden, Jeffry S., Thomas J. Zimmerman and David M. Worthen. "Hand-held Tonometers. An Evaluation by Medical Students." *Archives of Ophthalmology*. Vol. 89(2): pp. 110-112, February, 1973.
- Kaufman, Herbert E.. "Pressure Measurement: Which Tonometer?" *Investigative Ophthalmology*. Vol. 2: pp. 80-85. February, 1972.
- Kobayashi, A. S., L.-Y. S. Woo, C. Lawrence and W. A. Schlegel. "Analysis of the Corneo-scleral Shell by the Method of Direct Stiffness." *Journal of Biomechanics*. Vol. 4(5): pp. 323-330. October, 1971.
- Koetting, Robert A.. "Corneal Rigidity As A Factor in Contact Lens Wear." *Contacto*. Vol. 16(1): pp. 51-55. March, 1972.
- Kraus, H.. "(Tonometry in Infants. I. Differential Tonometry)." *Ceskoslovenska Oftalmologie*. Vol. 29(5): pp. 354-357. September, 1973.
- Kriegelstein, Guenter K. and Wolfgang K. Waller. "Goldmann Applanation Versus Hand-applanation and Schiotz Indentation Tonometry." *Albrecht Von Graefes Archiv fur Klinische und Experimentelle Ophthalmologie*. Vol. 194(1): pp. 11-16. March 3, 1975.

Langham, Maurice E., Wolfgang Leydhecker, Guenter Krieglstein and Wolfgang Waller. "Pneumatographic Studies on Normal and Glaucomatous Eyes." *Advances in Ophthalmology*. Vol. 32: pp. 108-133. 1976.

Levene, R. Z., "Glaucoma." *Archives of Ophthalmology*. Vol. 85: pp. 227-251.

Lyon, Catherine, W. K. McEwen and Marvin D. Shepherd. "Ocular Rigidity and Decay Curves Analyzed by Two Nonlinear Systems." *Investigative Ophthalmology*. Vol. 9(12): pp. 935-945. December, 1970.

Mc Ewen, W. K.. "An Advanced Method of Information Processing for Clinical Schiottz Tonograms." *Investigative Ophthalmology*. Vol. 12(11): pp.834-838. November, 1973.

Millodot, M.. "Effect of Corneal Curvature on Tonomat Measurements." *American Journal of Optometry*. Vol. 51: pp.12-14. January, 1974.

Moses, R.A.. "The Goldmann Applanation Tonometer." *American Journal of Ophthalmology*. Vol. 46: p. 865. 1958.

Moses, R. A.. "Theory of the Schiottz Tonometer and its Empirical Calibration." *American Ophthalmological Society Transcripts*. Vol. 69: pp. 494-562. 1971.

Moses, R. A., and W. J. Grodzki. "Theory and Calibration of the Schiottz Tonometer: II. The Resistance of the Cornea to Deformation." *Investigative Ophthalmology*. Vol. 10(7): pp. 534-543. July, 1971.

Moses, R.A., Editor. *Adler's Physiology of the Eye*. Chapter 6. "Intraocular Pressure." pp. 179-209. The C. V. Mosby Company, Saint Louis, 1975.

Nesterov, A. P. and M. B. Vurgaft. "(About Calibrometry of Filatov-Kalfa's Elastotonometer)." Oftal'mologicheskii Zhurnal. Vol. 30(1): pp. 46-47. 1975.

Özcetin, H. "The Coefficient of Ocular Rigidity." Turkish Oftal. Gazette. Vol. 4(2): pp. 70-77. 1974.

Poinoosawmy, D. and J. A. Roth. "Variations in Visual Acuity, Refraction and Corneal Curvature with Changes in Applanation Tension." British Journal of Ophthalmology. Vol. 58: pp. 523-528. May, 1974.

Previte, Louis R., James E. Burris, Vincent Carter and Bernard Carol. "Halberg Hand Applanation Tonometer: Statistical Analysis of Tonometry with Goldmann, Halberg and Schiotz Tonometers on 2,000 Eyes." Annals of Ophthalmology. Vol. 6(6): pp. 567-570. June, 1974.

Sagan, W. and K. N. Schwaderer. "Clinical Comparison of the Non-Contact Tonometer with the Mackay-Marg Tonometer." Journal of the American Optometric Association. Vol. 46: pp. 26-29. January, 1975.

Schapero, Max, David Cline and Henry W. Hofstetter. Dictionary of Visual Science. Second Edition. Chilton Book Company. Radnor, Pennsylvania. 1968

Scott, George Alan, Herman L. Tacker and Gerald P. Birkman. "Clinical Comparison of Four Tonometers." American Optometric Association Journal. Vol. 42(4): pp. 346-350. 1971.

Smith, J. L.. "Applanation Versus Schiotz Tonometry. The Syndrome of "Reverse Ocular Rigidity"." American Journal of Ophthalmology. Vol. 58: pp. 807-812. 1964.

Smith, J.L. et. al. "The Incidence of Schiotz-Appplanation Disparity." Archives of Ophthalmology. Vol. 77: pp.305-308. 1967.

Stelzer, R.. "The Relationship of Corneal Applanation to the Intra-Ocular Pressure in Enucleated Human Eyes." Albrecht Von Graefes Archiv fur Klinische und Experimentelle Ophthalmologie. Vol. 189(1): pp. 33-41. 1974.

Stepanik, Joseph. "The Mackay-Marg Tonometer. Correlation of the Tonogram to the Corneal Applanations Induced by the Tonometer." Acta Ophthalmologica. Vol. 48(6): pp. 1140-1144. 1970.

Stepanik, Joseph. "Das Mackay-Marg Tonometer. VI. Hornhaut-rigiditat und Augendruck ("The Mackay-Marg Tonometer. VI. Corneal Rigidity and Intraocular Pressure.") Albrecht Von Graefes Archiv fur Klinische und Experimentelle Ophthalmologie. Vol. 188(3): pp. 211-216. 1973.

Stepanik, Joseph. "Comparison of Modern Tonometers." Klinische Monatsblätter fur Augenheilkunde und Augenartzliche Fortbildung. Vol. 164(3): pp. 426-430. 1974.

Stepanik, Joseph. "Bulbushullen und Augendruck (The Coats of the Eyeball and the Intraocular Pressure.)" Klinische Monatsblätter fur Augenheilkunde und Augenartzliche Fortbildung. Vol. 166(1): pp.5-10. January, 1975.

Stuckey, G. C.. "Tonometry and the Imbert-Fick Law." Australian Journal of Ophthalmology. Vol. 2: pp.68-70. June, 1974.

- Stuckey, G. C.. "Tonometry-the Absolute Determination of the Original Intraocular Pressure." Australian Journal of Ophthalmology. Vol. 2: pp. 154-155. October, 1974.
- Syrdalen, Per. " Intraocular Pressure and Ocular Rigidity in Patients with Retinal Detachment. II. Postoperative Study." Acta Ophthalmologica. Vol. 48(5): pp. 1024-1044. 1970.
- Vucicevic, Zarko M. and Joanna Ralston. "Influences of the Volume and Hydration Changes on Scleral Rigidity." Annals of Ophthalmology. Vol. 4(9): pp. 715-732. September, 1972.
- Yamada, Eiichi. "On the Schiotz Tonometer- Comparison with the Applanation Tonometer." Acta Societatis Ophthalmologicae. Vol. 76(2): pp. 84-89. February, 1972.